

***AKARI* Near Infrared Spectroscopy: Detection of H₂O and CO₂ Ices toward Young Stellar Objects in the Large Magellanic Cloud**

Takashi SHIMONISHI^{1,2}, Takashi ONAKA¹, Daisuke KATO¹, Itsuki SAKON¹,
Yoshifusa ITA³, Akiko KAWAMURA⁴, and Hidehiro KANEDA⁵

ABSTRACT

We present the first results of *AKARI* Infrared Camera near-infrared spectroscopic survey of the Large Magellanic Cloud (LMC). We detected absorption features of the H₂O ice 3.05 μ m and the CO₂ ice 4.27 μ m stretching mode toward seven massive young stellar objects (YSOs). These samples are for the first time spectroscopically confirmed to be YSOs. We used a curve-of-growth method to evaluate the column densities of the ices and derived the CO₂/H₂O ratio to be 0.45 ± 0.17 . This is clearly higher than that seen in Galactic massive YSOs (0.17 ± 0.03). We suggest that the strong ultraviolet radiation field and/or the high dust temperature in the LMC may be responsible for the observed high CO₂ ice abundance.

Subject headings: circumstellar matter — stars: pre-main sequence — ISM: abundances — dust, extinction — ISM: molecules — Magellanic Clouds

1. Introduction

Properties of extragalactic young stellar objects (YSOs) provide us important information on the understanding of the diversity of YSOs in different galactic environments. The Large Magellanic Cloud (LMC), the nearest irregular galaxy to our Galaxy (~ 50 kpc; Alves

¹Department of Astronomy, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0003, Japan

²shimonishi@astron.s.u-tokyo.ac.jp

³National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁴Department of Astrophysics, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan

⁵Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan

2004), offers an ideal environment for this study since it holds a unique metal-poor environment (Luck et al. 1998). Because of its proximity and nearly face-on geometry, various types of surveys have been performed toward the LMC (e.g., Zaritsky et al. 2004; Meixner et al. 2006; Kato et al. 2007, and references therein).

An infrared spectrum of YSOs shows absorption features of various ices which are thought to be an important reservoir of heavy elements and complex molecules in a cold environment such as a dense molecular cloud or an envelope of a YSO (e.g., Chiar et al. 1998; Nummelin et al. 2001; Whittet et al. 2007; Boogert et al. 2008). These ices are thought to be taken into planets and comets as a result of subsequent planetary formation activity (Ehrenfreund & Schutte 2000). Studying the compositions of ices as functions of physical environments is crucial to understand the chemical evolution in circumstellar environments of YSOs and is a key topic of astrophysics. H_2O and CO_2 ices are ubiquitous and are major components of interstellar ices (van Dishoeck & Blake 1998; Boogert & Ehrenfreund 2004). Since the absorption profile of the ices is sensitive to a chemical composition of icy grain mantles and a thermal history of local environments, the ices are important tracers to investigate the properties of YSOs. However, our knowledge about the ices around extragalactic YSOs is limited because few observations have been performed toward extragalactic YSOs. Therefore infrared spectroscopic observations toward YSOs in the LMC are important if we are to improve our understanding of the influence of galactic environments on the properties of YSOs and ices.

AKARI is the first Japanese satellite dedicated to an infrared astronomy launched in February 2006 (Murakami et al. 2007). We have performed a near infrared spectroscopic survey of the LMC using a powerful spectroscopic survey capability of Infrared Camera (IRC; Onaka et al. 2007) on board *AKARI*. In this letter, we present $2.5\text{--}5\mu\text{m}$ spectra of newly confirmed YSOs in the LMC with our survey, and discuss the abundance of H_2O and CO_2 ice.

2. Observations and Data Reduction

The observations reported here were obtained as a part of the *AKARI* IRC survey of the LMC (Ita et al. 2008). An unbiased slitless prism spectroscopic survey of the LMC has been performed since May 2006. In this survey, the IRC02b *AKARI* astronomical observing template (AOT) with the NP spectroscopy mode was used to obtain low resolution spectra ($R \sim 20$) between 2.5 and $5\mu\text{m}$. By June 2008, 621 pointing observations were performed toward the LMC, and about 10 deg^2 area was observed. The near-infrared (NIR) spectroscopic survey is continuing even after the exhaustion of the liquid helium in August 2008.

The spectral analysis was performed using the standard IDL package prepared for the reduction of *AKARI* IRC spectra (Ohya et al. 2007). Raw data were converted to dark-subtracted, linearity-corrected, and flat-field-corrected frames. Three pixels were integrated in the spatial direction to extract a point-source spectrum from the slitless spectroscopic image. Background levels of each source were estimated from the signal counts of the adjacent regions on both sides of the target spectrum. The wavelength calibration accuracy is estimated to be about $\sim 0.01\mu\text{m}$ (Ohya et al. 2007)

3. The selection of YSOs

We select infrared-bright objects from the point-source catalog of the *Spitzer* SAGE project (Meixner et al. 2006) with the following selection criteria: (1) $[3.6] - [4.5] > 0.3$ and $[5.8] - [8.0] > 0.6$, and (2) $[3.6] < 12$ and $[4.5] < 11.5$, where [wavelength] represents the photometric value in magnitude at each wavelength in microns. Criterion (1) refers to the YSO model of Whitney et al. (2004), and criterion (2) comes from the detection limit of the *AKARI* IRC NP mode. This rough selection is applied to the sources located in the survey area of *AKARI* IRC, and about 300 sources are selected. These photometrically selected sources include not only massive YSOs, but also a large number of dusty evolved stars since their infrared spectral energy distributions (SEDs) are similar to each other. For the accurate selection of YSOs, we select the sources that show absorption features of the $3.05\mu\text{m}$ H_2O ice and the $4.27\mu\text{m}$ CO_2 ice stretching mode in their NIR spectra taken by the present spectroscopic survey. The presence of CO_2 ice is strong evidence of YSOs since the detection of CO_2 ice toward dusty evolved stars has not been reported (Barlow 1999; Sylvester et al. 1999). Spectral overlapping with other sources located in a dispersion direction is a serious problem for slit-less spectroscopy, which makes it difficult to obtain reliable spectra. We check the overlapping contamination by visual inspection, and we only use the sources without such contamination in the following analysis.

As a result, we spectroscopically confirmed seven massive YSOs in the LMC for the first time. The sources are listed in Table 1 with the observation parameters. Six of the seven sources are included in the recent YSO candidates catalog (Whitney et al. 2008), and one source (ST6) is a newly found YSO. The spectra of these sources are shown in Fig 1 together with the results of spectral fitting (see §4 for details). The absorption features of H_2O and CO_2 ices are rather broadened due to the low spectral resolution of the *AKARI* IRC NP spectroscopy mode but clearly detected. This is the first clear detection of the $4.27\mu\text{m}$ stretching mode of CO_2 ice toward extragalactic YSOs. In addition, unresolved emission of PAHs and the hydrogen recombination line $\text{Pf}\delta$ around $3.3\mu\text{m}$, the $\text{Br}\alpha$ line at $4.05\mu\text{m}$

and blended absorption features of $4.62\ \mu\text{m}$ XCN and $4.67\ \mu\text{m}$ CO ices around $4.65\ \mu\text{m}$ are detected toward several sources. However, it is difficult to evaluate the column densities of XCN and CO ices accurately with the present low spectral resolution data.

4. Spectral Fitting

We fit a polynomial of the second to fourth order to the continuum regions and divide the spectra by the fitted continuum (Fig 1). The wavelength regions for the continuum are set to be $2.5\text{--}2.7\ \mu\text{m}$, $3.6\text{--}3.7\ \mu\text{m}$, $4.0\text{--}4.15\ \mu\text{m}$, and $4.9\text{--}5.0\ \mu\text{m}$ (Gibb et al. 2004).

Due to the low spectral resolution of the IRC NP spectroscopy mode, direct comparison of the observed spectra with the absorption profiles of laboratory ices is difficult. However, the equivalent width of absorption does not depend on the spectral resolution of the spectrum. Therefore we used a curve-of-growth method to derive the column densities of the ices. A Gaussian profile with the fixed central wavelength is fitted to the absorption bands to derive the equivalent width (Fig 1). The vertical axis of the plot is shown in units of the normalized flux because it is difficult to estimate the optical depth directly from the present low resolution data. We use the laboratory absorption profiles of H_2O and CO_2 ices taken from the Leiden Molecular Astrophysics database (Ehrenfreund et al. 1996) to calculate the curve of growth. The profiles of pure H_2O ice and mixture of $\text{H}_2\text{O} : \text{CO}_2$ (100:14) ice both at 10K are used for the calculation since these compositions are typical in the interstellar ices (Nummelin et al. 2001; Gibb et al. 2004). The present spectrum cannot resolve the polar and apolar CO_2 ice features, and the present analysis assumes the polar CO_2 ice only. However, contribution of the apolar ice is generally small toward YSOs (Gerakines et al. 1995, 1999). The FWHM of the absorption profiles of H_2O and CO_2 ices change by approximately 10 % depending on the compositions of the ices (Gerakines et al. 1999; Gibb et al. 2004). We confirm that this range of the variation in the FWHM makes negligible effects on the derived column densities compared with the observational errors. We adopt the band strengths of H_2O and CO_2 ices to be 2.0×10^{-16} and $7.6 \times 10^{-17}\ \text{cm molecule}^{-1}$ (Gerakines et al. 1995), respectively. The derived column densities are listed in Table 1.

Since this study applies the curve of growth method to the calculation of the ice column density for the first time, we check the validity of our method. We use a few Galactic YSO spectra taken by *ISO* SWS whose column density of H_2O and CO_2 ices are derived in Gibb et al. (2004). These high-resolution spectra are converted to low-resolution spectra by convolving the slit function of the NP spectroscopy mode, and then we derive the column density for these converted spectra using the same method described above. The comparison of the obtained column density with the value presented in Gibb et al. (2004) shows that

the differences are within 10 % and 20 % for the H₂O ice and the CO₂ ice, respectively. The result plotted in Fig 2 includes this uncertainty in addition to the observational error.

5. Results and Discussion

The obtained column densities of H₂O and CO₂ ices are plotted in Fig 2. The error bars become larger for the larger column density due to the saturation effect of the curve-of-growth. A linear fit to the data points indicates that the CO₂/H₂O ice column density ratio in the LMC is 0.45 ± 0.17 . The large uncertainty mainly comes from the errors in the curve-of-growth analysis. For comparison, column densities of Galactic massive YSOs taken from Gibb et al. (2004) and their CO₂/H₂O ice column density ratio of 0.17 ± 0.03 (Gerakines et al. 1999) are also plotted in Fig 2. A similar CO₂/H₂O ratio of 0.18 ± 0.04 is also observed toward a Galactic quiescent dark cloud (Whittet et al. 2007), while a relatively high CO₂/H₂O ratio of 0.32 ± 0.02 is observed toward Galactic low- and intermediate- mass YSOs, and some of them reach ~ 0.4 (Pontoppidan et al. 2008). Although the uncertainty is large, it is clear from the present results that the CO₂/H₂O ice ratio in the LMC is higher than the typical ratios of the Galactic objects. Since the distribution range of the H₂O ice column density in the LMC is comparable to that of the massive Galactic YSOs, it can be concluded that the abundance of the CO₂ ice is higher in the LMC. The present results suggest that the different galactic environment of the LMC is responsible for the high CO₂ abundance.

The formation mechanism of CO₂ ice in circumstellar environments of YSOs is not understood, however a number of scenarios have been proposed. Several laboratory experiments indicate that the CO₂ ice is efficiently produced by UV photon irradiation to H₂O-CO binary ice mixtures (e.g., Watanabe et al. 2007). The LMC has an order-of-magnitude stronger UV radiation field than our Galaxy due to its active massive star formation (Israel & de Graauw 1986), which could lead to the higher CO₂/H₂O ratio in the LMC. The high CO₂/H₂O ratio toward a YSO in the LMC is also reported in van Loon et al. (2005), and they suggest that a different radiation environment in the LMC is one of the reasons for the high CO₂ abundance. On the other hand, the model of diffusive surface chemistry suggests that high abundance of CO₂ ice can be produced at relatively high dust temperatures (Bergin, Neufeld, & Melnick 1999; Ruffle & Herbst 2001). Several studies have reported that the dust temperature in the LMC is generally higher than in our Galaxy based on far-infrared to submillimeter observations of diffuse emission (e.g., Aguirre et al. 2003; Sakon et al. 2006). Therefore the high dust temperature may also have an effect on the high CO₂ ice abundance in the LMC.

It is difficult to separate the effect of the UV radiation field and the dust temperature on

the high abundance of CO₂ ice in the LMC by our low-resolution NIR spectra. The 4.62 μm XCN feature is known to be indicative of strong UV irradiation (Bernstein, Sandford, & Allamandola 2000; Spoon et al. 2003). On the other hand, detailed profile analysis of the 3.05 μm H₂O ice stretching mode and the 15.2 μm CO₂ ice bending mode should reveal the temperature and compositions of the ices (Ehrenfreund et al. 1996; Oberg et al. 2007). Future observations of the XCN feature and the H₂O and CO₂ ice features with a sufficient wavelength resolution will be useful to investigate this problem.

6. Summary

We performed a near-infrared spectroscopic survey of the LMC with *AKARI* IRC. We spectroscopically confirmed seven massive YSOs that show absorption features of H₂O and CO₂ ices. This is the first detection of the 4.27 μm CO₂ ice feature toward extragalactic YSOs. The derived ice column densities indicate that the abundance of CO₂ ice is clearly higher in the LMC than our Galaxy. The relatively strong UV radiation field and/or high dust temperature in the LMC may be responsible for the observed high abundance of CO₂ ice. Our study shows the difference in the chemical composition around extragalactic YSOs, suggesting that extragalactic YSOs hold quite different environments from Galactic ones.

AKARI is a JAXA project with the participation of ESA. We thank all the members of the *AKARI* project for their continuous help and support. We also thank Youichi Ohyama for great support in the analysis of spectroscopic data, and we thank the *AKARI* IRC team and the LMC team for their continuous support. This work is supported by a Grant-in-Aid for Scientific Research from the JSPS.

REFERENCES

- Aguirre, J. E., et al., 2003, *ApJ*, 596, 273
- Alves, D. R., 2004, *New A Rev.*, Vol.48, Issue 9, p.659-665
- Barlow, M. J., 1999, *I.A.U. Symposium* 191, 353
- Bergin, E. A., Neufeld, D. A., & Melnick, G. J., 1999, *ApJ*, 510, 145
- Bernstein, M. P., Sandford, S. A., & Allamandola, L., J. 2000, *ApJ*, 542, 892
- Boogert, A. C. A., & Ehrenfreund, P., 2004, *ASP Conference Series*, Vol.309

- Boogert, A. C. A., et al., 2008, *ApJ*, 678, 985
- Chiar, J. E., Gerakines, P. A., Whittet, D. C. B., Pendleton, Y. J., Tielens, A. G. G. M., Adamson, A. J., & Boogert A. C. A., 1998, *ApJ*, 498, 716
- Ehrenfreund, P., Boogert, A. C. A., Gerakines, P. A., Jansen, D. J., Schutte, W. A., Tielens, A. G. G. M., & van Dishoeck, E. F., 1996, *A&A*, 315, L341
- Ehrenfreund, P., & Schutte, W. A., 2000, *Adv. Space Res.*, Vol.25, No.11, pp.2177-2188
- Gerakines, P. A., Schutte, W. A., Greenberg, J. M., & van Dishoeck, E. F. 1995, *A&A*, 296, 810
- Gerakines, P. A., et al., 1999, *ApJ*, 522, 357
- Gibb, E. L., Whittet, D. C. B., Schutte, W. A., et al., 2000, *ApJ*, 536, 347
- Gibb, E. L., Whittet, D. C. B., Boogert, A. C. A., & Tielens, A. G. G. M., 2004, *ApJS*, 151, 35
- Israel, F. P., & de Graauw, Th., van de Stadt, D., & de Vries, C. P., 1986, *ApJ*, 303, 186
- Ita, Y., et al., *PASJ*, submitted
- Kato, D., et al., 2007, *PASJ*, 59, 615
- Luck, R. E., Moeffett, T. J., Barnes III, T. G., & Gieren, P. W., 1998, *AJ*, 115, 605
- Meixner, M., et al., 2006, *AJ*, 132, 2268
- Murakami, H. et al., 2007, *PASJ*, 59, S369
- Nummelin, A., Whittet, D. C. B., Gibb, E. L., Gerakines, P. A., & Chiar, J. E., 2001, *ApJ*, 558, 185
- Oberg, Fraser, H. J., Boogert, A. C. A., Bisschop, S. E., Fuchs, G. W., van Dishoeck, E. F., & Linnartz, H., 2007, *A&A*, 462, 1187
- Ohyama, Y. et al., 2007, *PASJ*, 59, S411
- Onaka, T. et al. 2007, *PASJ*, 59, S401
- Pontoppidan, K. M., et al., 2008, *ApJ*, 678, 1005
- Ruffle D. P., & Herbst E., 2001, *MNRAS*, 324, 1054

- Sakon, I. et al., 2006, *ApJ*, 651, 174
- Spoon, H. W. W., Moorwood, A. F. M., Pontoppidan, K. M., Cami, J., Kregel, M., Lutz, D., & Tielens, A. G. G. M., 2003, *A&A*, 402, 499
- Sylvester, R. J., Kemper, F., Barlow, M. J., de Jong, T., Waters, L. B. F. M., Tielens, A. G. G. M., & Omont, A., 1999, *A&A*, 352, 587
- van Dishoeck, E. F. & Blake, G. A., 1998, *ARA&A*, 36, 317
- van Loon, J. Th., et al., 2005, *MNRAS*, 364, L71
- Watanabe, N., Mouri, O., Nagaoka, A., Chigai, T., Kouchi, A., & Pirronello, V., 2007, *ApJ*, 668, 1001
- Whitney, B. A., Indebetouw, R., Bjorkman, J. E., & Wood, K., 2004, *ApJ*, 617, 1177
- Whitney, B. A., et al., 2008, *AJ*, 136, 18
- Whittet, D. C. B., Shenoy, S. S., Bergin, E. A., Chiar, J. E., Gerakines, P. A., Gibb, E. L., Melnick, G. J., & Neufeld, D. A., 2007, *ApJ*, 655, 332
- Zaritsky, D., Harris, J., Thompson, I. B., & Grebel, E. K., 2004, *AJ*, 128, 1606

Table 1. OBSERVATION PARAMETERS AND COLUMN DENSITIES OF ICES

Number	AKARI ID	Obs. ID	Obs. Date	Other Name	RA[J2000]	DEC[J2000]	N(H ₂ O) (10 ¹⁷ cm ⁻²)	N(CO ₂) ^a (10 ¹⁷ cm ⁻²)
ST1	J053931-701216	2211375	2007 Apr 12	05393117-7012166 ^a	05:39:31.15	-70:12:16.8	9.6 ^{+1.9} _{-1.9}	6.7 ^{+4.8} _{-3.6}
ST2	J052212-675832	2210229	2006 Nov 24	NGC 1936	05:22:12.56	-67:58:32.2	11.1 ^{+1.6} _{-1.4}	3.1 ^{+1.7} _{-1.5}
ST3	J052546-661411	2213132	2007 Jun 22	 ^b	05:25:46.69	-66:14:11.3	29.7 ^{+4.5} _{-4.6}	15.5 ^{+13.8} _{-9.4}
ST4	J051449-671221	2200073	2006 Jun 6	IRAS F05148-6715	05:14:49.41	-67:12:21.5	18.7 ^{+2.3} _{-2.3}	8.2 ^{+4.5} _{-3.7}
ST5	J053054-683428	2213061	2007 Mar 13	IRAS 05311-6836	05:30:54.27	-68:34:28.2	31.7 ^{+4.7} _{-4.5}	12.4 ^{+9.1} _{-6.8}
ST6	J053941-692916	2201137	2006 Oct 25	05394112-6929166 ^a	05:39:41.08	-69:29:16.8	59.1 ^{+38.4} _{-25.0}	...
ST7	J052351-680712	2210220	2006 Nov 29	IRAS05240-6809	05:23:51.15	-68:07:12.2	...	55.3 ^{+40.3} _{-30.3}

^a2MASS ID

^bThe source is in a cluster




Fig. 1.— *AKARI* IRC 2.5–5 μ m spectra of YSOs in the LMC. Left: Plots of $\log_{10}$ flux(Jy) vs. wavelength(μ m). Open circles represent the points that used for the continuum determination. Dashed lines represent derived continuum. Right: Plots of normalized flux ($1 - F/F_c$; F and F_c represent observed flux and continuum flux, respectively) vs. wavelength(μ m). Shaded regions around 3.0 μ m and 4.3 μ m represent fitted Gaussian for absorption features of H₂O and CO₂ ices, and the areas correspond to the equivalent width. The positions of H₂O, CO₂, XCN, CO ice absorption bands, PAH emission bands, and hydrogen recombination lines are shown.




Fig. 2.— CO₂ ice vs. H₂O ice column density in a unit of 10¹⁷ cm⁻². Open squares with error bars represent the results of this study. Filled squares represent those of Galactic massive YSOs (Gibb et al. 2004). Upper and lower solid lines represent CO₂/H₂O $\sim$ 0.45 and 0.17, respectively. The sources ST6 and ST7 are not plotted due to their large errors.